

Animal Medical Center of Troy
994 West South Blvd. at Livernois
Rochester Hills, MI 48307-4937
(248) 852-9800
www.amctroy.com

Poppy

1 year old
13 lbs.

Beth Widdows Westie Rescue
1371 W. South Blvd.
Troy, MI 48098



02/27/2015 Theodoroff **(Note)** Final Vaccine series, final Ivermectin injection
Final Vaccine series, final Ivermectin .13 ml injection (lung worms seen with Lily)
3 months of Sentinel Spectrum
Doing very well, Very sweet puppy

DHLPP vaccine

This vaccine protects against distemper, hepatitis, leptospirosis, parvovirus and parainfluenza.

Distemper is an easily transmitted viral disease of dogs that is usually fatal in puppies and young dogs without immunity. Symptoms include diarrhea, discharge from the eyes and nose, fever, anorexia, and respiratory difficulties. The disease can also cause neurological symptoms such as seizures. **Parainfluenza** is a viral disease and is considered a minor respiratory infection among healthy adult dogs. It can be severe in puppies or debilitated dogs, and is characterized by mild fever, nasal discharge, reddened tonsils and a harsh, non-productive cough. The two strains of adenovirus cause infectious **hepatitis** and respiratory infection. Most adult dogs can recover from hepatitis, but puppies often can't. Adenovirus type-1 can cause fever, diarrhea and severe liver, kidney and eye damage. Adenovirus type-2 can contribute to respiratory disease (canine cough).

Parvovirus is a highly contagious disease often resulting in death, especially in puppies. Parvovirus normally infects the dog's intestines, lymphoid tissue and bone marrow. The result is vomiting; severe, often fatal, bloody diarrhea; and severe dehydration. It mostly affects young pets that have not been adequately immunized. **Leptospirosis** is a bacterial disease that causes permanent kidney and liver damage. It can be fatal and is contagious to other animals and to humans.

This vaccine is recommended for all puppies and dogs. Puppies should be vaccinated every 3 weeks between 6 and 16 weeks of age. Dogs should be vaccinated at 1 year and 2 years of age, then every 3 years thereafter. The leptospirosis component is boosted every year.

This vaccine was given under the skin over the left shoulder. Dogs may be lethargic or sore for 24 hours following vaccination. Facial swelling, hives, vomiting and diarrhea could be signs of a serious vaccine reaction and should be reported to a veterinarian immediately.

The brand of vaccine given today is Vanguard Plus 5 L4 made by Zoetis.

Next vaccine 1 year from today

Bordetella vaccine, injectable

Tracheobronchitis ("Kennel Cough") is an easily transmitted upper respiratory disease with a harsh, dry cough. It can be mild or very serious, sometimes leading to pneumonia. This vaccine protects against bordetella bronchiseptica, one of the most important causes of tracheobronchitis. Tracheobronchitis is commonly acquired by dogs that come into contact with groups of other dogs at kennels, dog parks, day care or grooming facilities. This vaccine is recommended for all puppies and dogs.

This vaccine was given under the skin in the neck area. Dogs may be lethargic or sore for 24 hours following vaccination. Facial swelling, hives, vomiting and diarrhea could be signs of a serious vaccine reaction and should be reported to a veterinarian immediately.

The brand of vaccine given today is Bronchicine CAe made by Zoetis.

Next vaccine 1 year from today

Sentinel Spectrum (milbemycin oxime/lufenuron/praziquantel)

Sentinel Spectrum can both **prevent and treat** intestinal **roundworm, hookworm, whipworm, and tapeworm infections**. These parasites can cause diarrhea in your dog, and can be passed to humans, often with very serious consequences. Your dog is exposed to these parasites all year round. Sentinel Spectrum also **prevents heartworm**, a sometimes-fatal disease of dogs that is carried by mosquitos. Sentinel spectrum **prevents flea infestations** by keeping flea eggs from hatching and flea larvae from maturing. Side effects are extremely rare. Please call your veterinarian if your dog experiences vomiting, diarrhea, lethargy, itching, drooling, weakness or seizures.

Give 1 chewable tablet by mouth on the same day each month. Give every month - all year round. Give this tablet with a meal.

These chews are highly flavored, please keep unused chews in an area that your pet can not access.

BRAVECTO™ - for tick control

Bravecto is a tasty oral chew that gives your dog up to 12 weeks of protection against fleas and ticks. It provides broad-spectrum and long-lasting protection which controls 4 tick species. (98% effective after 12 hours for 12 weeks, Lone Star Tick for 8 weeks). Being an oral medication, it can't be rubbed or washed off. There is no chemical exposure to the family. Bravecto will also kill adult fleas for 12 weeks.

The active ingredient in BRAVECTO, Fluralaner, has been shown to be safe at 5 times the recommended dose. There are no known contraindications for the use of the medication. Bravecto can be used in adult dogs, and it's even safe for use in breeding, pregnant, and lactating dogs. For use in growing puppies, they should be 6 months of age or older and weigh at least 4.4 lbs.

For any questions on Poppy's health, please call the clinic.